

# RT<sup>2</sup> Profiler PCR Array (Rotor-Gene<sup>®</sup> Format)

## Mouse Innate & Adaptive Immune Responses

Cat. no. 330231 PAMM-052ZR

For pathway expression analysis

| Format                                       | For use with the following real-time cyclers |
|----------------------------------------------|----------------------------------------------|
| RT <sup>2</sup> Profiler PCR Array, Format R | Rotor-Gene Q, other Rotor-Gene cyclers       |

### Description

The Mouse Innate & Adaptive Immune Responses RT<sup>2</sup> Profiler PCR Array profiles the expression of 84 genes involved in the host response to bacterial infection and sepsis. This array includes genes related to the IL-1R and Toll-like Receptor (TLR) Signaling Pathways including IL-1R and TLR genes involved in the detection of pathogens. Genes related to the host defense to bacteria are represented on this array including genes involved in the detection of bacteria, and genes involved in the acute-phase response, complement activation, the inflammatory response, and the antibacterial humoral response. Genes involved in the innate immune response and septic shock are also included on this array. Using real-time PCR, you can easily and reliably analyze expression of a focused panel of genes related to the innate and adaptive immune responses with this array.

For further details, consult the *RT<sup>2</sup> Profiler PCR Array Handbook*.

### Shipping and storage

RT<sup>2</sup> Profiler PCR Arrays in the Rotor-Gene format are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products.

For long term storage, keep plates at –20°C.

**Note:** Ensure that you have the correct RT<sup>2</sup> Profiler PCR Array format for your real-time cycler (see table above).

**Note:** Open the package and store the products appropriately immediately on receipt.



## Array layout

The 96 real-time assays in the Rotor-Gene format are located in wells 1–96 of the Rotor-Disc™ (plate A1–A12=Rotor-Disc 1–12, plate B1–B12=Rotor-Disc 13–24, etc.). To maintain data analysis compatibility, wells 97–100 do not contain real-time assays but will contain master mix to account for weight balance.

## Gene table: RT<sup>2</sup> Profiler PCR Array

| Position | UniGene   | GenBank      | Symbol | Description                                          |
|----------|-----------|--------------|--------|------------------------------------------------------|
| A01      | Mm.330510 | NM_011318    | Apcs   | Serum amyloid P-component                            |
| A02      | Mm.19131  | NM_009778    | C3     | Complement component 3                               |
| A03      | Mm.247623 | NM_007577    | C5ar1  | Complement component 5a receptor 1                   |
| A04      | Mm.1051   | NM_009807    | Casp1  | Caspase 1                                            |
| A05      | Mm.867    | NM_011331    | Ccl12  | Chemokine (C-C motif) ligand 12                      |
| A06      | Mm.284248 | NM_013653    | Ccl5   | Chemokine (C-C motif) ligand 5                       |
| A07      | Mm.1337   | NM_009916    | Ccr4   | Chemokine (C-C motif) receptor 4                     |
| A08      | Mm.14302  | NM_009917    | Ccr5   | Chemokine (C-C motif) receptor 5                     |
| A09      | Mm.8007   | NM_009835    | Ccr6   | Chemokine (C-C motif) receptor 6                     |
| A10      | Mm.442098 | NM_007720    | Ccr8   | Chemokine (C-C motif) receptor 8                     |
| A11      | Mm.3460   | NM_009841    | Cd14   | CD14 antigen                                         |
| A12      | Mm.2209   | NM_013488    | Cd4    | CD4 antigen                                          |
| B01      | Mm.271833 | NM_011611    | Cd40   | CD40 antigen                                         |
| B02      | Mm.4861   | NM_011616    | Cd40lg | CD40 ligand                                          |
| B03      | Mm.89474  | NM_009855    | Cd80   | CD80 antigen                                         |
| B04      | Mm.1452   | NM_019388    | Cd86   | CD86 antigen                                         |
| B05      | Mm.1858   | NM_001081110 | Cd8a   | CD8 antigen, alpha chain                             |
| B06      | Mm.28767  | NM_007768    | Crp    | C-reactive protein, pentraxin-related                |
| B07      | Mm.4922   | NM_009969    | Csf2   | Colony stimulating factor 2 (granulocyte-macrophage) |
| B08      | Mm.877    | NM_021274    | Cxcl10 | Chemokine (C-X-C motif) ligand 10                    |
| B09      | Mm.12876  | NM_009910    | Cxcr3  | Chemokine (C-X-C motif) receptor 3                   |
| B10      | Mm.86382  | NM_172689    | Ddx58  | DEAD (Asp-Glu-Ala-Asp) box polypeptide 58            |
| B11      | Mm.3355   | NM_010177    | Fas1   | Fas ligand (TNF superfamily, member 6)               |
| B12      | Mm.182291 | NM_054039    | Foxp3  | Forkhead box P3                                      |
| C01      | Mm.313866 | NM_008091    | Gata3  | GATA binding protein 3                               |
| C02      | Mm.88795  | NM_010391    | H2-Q10 | Histocompatibility 2, Q region locus 10              |
| C03      | Mm.439648 | NM_010398    | H2-T23 | Histocompatibility 2, T region locus 23              |
| C04      | Mm.435508 | NM_010493    | Icam1  | Intercellular adhesion molecule 1                    |
| C05      | Mm.14091  | NM_010503    | Ifna2  | Interferon alpha 2                                   |
| C06      | Mm.502    | NM_010508    | Ifnar1 | Interferon (alpha and beta) receptor 1               |
| C07      | Mm.1245   | NM_010510    | Ifnb1  | Interferon beta 1, fibroblast                        |
| C08      | Mm.240327 | NM_008337    | Ifng   | Interferon gamma                                     |
| C09      | Mm.549    | NM_010511    | Ifngr1 | Interferon gamma receptor 1                          |
| C10      | Mm.874    | NM_010548    | Il10   | Interleukin 10                                       |
| C11      | Mm.1284   | NM_008355    | Il13   | Interleukin 13                                       |
| C12      | Mm.5419   | NM_010552    | Il17a  | Interleukin 17A                                      |
| D01      | Mm.1410   | NM_008360    | Il18   | Interleukin 18                                       |
| D02      | Mm.15534  | NM_010554    | Il1a   | Interleukin 1 alpha                                  |
| D03      | Mm.222830 | NM_008361    | Il1b   | Interleukin 1 beta                                   |
| D04      | Mm.896    | NM_008362    | Il1r1  | Interleukin 1 receptor, type I                       |
| D05      | Mm.14190  | NM_008366    | Il2    | Interleukin 2                                        |
| D06      | Mm.125482 | NM_031252    | Il23a  | Interleukin 23, alpha subunit p19                    |
| D07      | Mm.276360 | NM_021283    | Il4    | Interleukin 4                                        |
| D08      | Mm.4461   | NM_010558    | Il5    | Interleukin 5                                        |
| D09      | Mm.1019   | NM_031168    | Il6    | Interleukin 6                                        |
| D10      | Mm.38241  | NM_008363    | Il6st  | Interleukin-6 receptor-associated kinase 1           |
| D11      | Mm.3960   | NM_016849    | Irf3   | Interferon regulatory factor 3                       |
| D12      | Mm.3233   | NM_016850    | Irf7   | Interferon regulatory factor 7                       |
| E01      | Mm.262106 | NM_008401    | Itgam  | Integrin alpha M                                     |
| E02      | Mm.275839 | NM_008413    | Jak2   | Janus kinase 2                                       |
| E03      | Mm.116844 | NM_016923    | Ly96   | Lymphocyte antigen 96                                |
| E04      | Mm.45436  | NM_017372    | Lyz2   | Lysozyme 2                                           |
| E05      | Mm.196581 | NM_011949    | Mapk1  | Mitogen-activated protein kinase 1                   |
| E06      | Mm.21495  | NM_016700    | Mapk8  | Mitogen-activated protein kinase 8                   |
| E07      | Mm.30045  | NM_010776    | Mbl2   | Mannose-binding lectin (protein C) 2                 |
| E08      | Mm.4668   | NM_010824    | Mpo    | Myeloperoxidase                                      |
| E09      | Mm.33996  | NM_010846    | Mx1    | Myxovirus (influenza virus) resistance 1             |

| Position | UniGene   | GenBank   | Symbol   | Description                                                                         |
|----------|-----------|-----------|----------|-------------------------------------------------------------------------------------|
| E10      | Mm.213003 | NM_010851 | Myd88    | Myeloid differentiation primary response gene 88                                    |
| E11      | Mm.256765 | NM_008689 | Nfkb1    | Nuclear factor of kappa light polypeptide gene enhancer in B-cells 1, p105          |
| E12      | Mm.170515 | NM_010907 | Nfkbia   | Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha |
| F01      | Mm.54174  | NM_145827 | Nlrp3    | NLR family, pyrin domain containing 3                                               |
| F02      | Mm.28498  | NM_172729 | Nod1     | Nucleotide-binding oligomerization domain containing 1                              |
| F03      | Mm.222633 | NM_145857 | Nod2     | Nucleotide-binding oligomerization domain containing 2                              |
| F04      | Mm.828    | NM_009019 | Rag1     | Recombination activating gene 1                                                     |
| F05      | Mm.4372   | NM_011281 | Rorc     | RAR-related orphan receptor gamma                                                   |
| F06      | Mm.2913   | NM_013612 | Slc11a1  | Solute carrier family 11 (proton-coupled divalent metal ion transporters), member 1 |
| F07      | Mm.277406 | NM_009283 | Stat1    | Signal transducer and activator of transcription 1                                  |
| F08      | Mm.249934 | NM_011486 | Stat3    | Signal transducer and activator of transcription 3                                  |
| F09      | Mm.1550   | NM_011487 | Stat4    | Signal transducer and activator of transcription 4                                  |
| F10      | Mm.121721 | NM_009284 | Stat6    | Signal transducer and activator of transcription 6                                  |
| F11      | Mm.477879 | NM_019507 | Tbx21    | T-box 21                                                                            |
| F12      | Mm.203952 | NM_174989 | Ticam1   | Toll-like receptor adaptor molecule 1                                               |
| G01      | Mm.273024 | NM_030682 | Tlr1     | Toll-like receptor 1                                                                |
| G02      | Mm.87596  | NM_011905 | Tlr2     | Toll-like receptor 2                                                                |
| G03      | Mm.33874  | NM_126166 | Tlr3     | Toll-like receptor 3                                                                |
| G04      | Mm.38049  | NM_021297 | Tlr4     | Toll-like receptor 4                                                                |
| G05      | Mm.116894 | NM_016928 | Tlr5     | Toll-like receptor 5                                                                |
| G06      | Mm.42146  | NM_011604 | Tlr6     | Toll-like receptor 6                                                                |
| G07      | Mm.23979  | NM_133211 | Tlr7     | Toll-like receptor 7                                                                |
| G08      | Mm.196676 | NM_133212 | Tlr8     | Toll-like receptor 8                                                                |
| G09      | Mm.44889  | NM_031178 | Tlr9     | Toll-like receptor 9                                                                |
| G10      | Mm.1293   | NM_013693 | Tnf      | Tumor necrosis factor                                                               |
| G11      | Mm.292729 | NM_009424 | Traf6    | Tnf receptor-associated factor 6                                                    |
| G12      | Mm.20249  | NM_018793 | Tyk2     | Tyrosine kinase 2                                                                   |
| H01      | Mm.328431 | NM_007393 | Actb     | Actin, beta                                                                         |
| H02      | Mm.163    | NM_009735 | B2m      | Beta-2 microglobulin                                                                |
| H03      | Mm.343110 | NM_008084 | Gapdh    | Glyceraldehyde-3-phosphate dehydrogenase                                            |
| H04      | Mm.3317   | NM_010368 | Gusb     | Glucuronidase, beta                                                                 |
| H05      | Mm.2180   | NM_008302 | Hsp90ab1 | Heat shock protein 90 alpha (cytosolic), class B member 1                           |
| H06      | N/A       | SA_00106  | MGDC     | Mouse Genomic DNA Contamination                                                     |
| H07      | N/A       | SA_00104  | RTC      | Reverse Transcription Control                                                       |
| H08      | N/A       | SA_00104  | RTC      | Reverse Transcription Control                                                       |
| H09      | N/A       | SA_00104  | RTC      | Reverse Transcription Control                                                       |
| H10      | N/A       | SA_00103  | PPC      | Positive PCR Control                                                                |
| H11      | N/A       | SA_00103  | PPC      | Positive PCR Control                                                                |
| H12      | N/A       | SA_00103  | PPC      | Positive PCR Control                                                                |

## Related products

For optimal performance, RT<sup>2</sup> Profiler PCR Arrays should be used together with the RT<sup>2</sup> First Strand Kit for cDNA synthesis and RT<sup>2</sup> SYBR<sup>®</sup> Green qPCR Mastermixes for PCR.

| Product                                             | Contents                                                                                                 | Cat. no. |
|-----------------------------------------------------|----------------------------------------------------------------------------------------------------------|----------|
| RT <sup>2</sup> First Strand Kit (12)               | Enzymes and reagents for cDNA synthesis                                                                  | 330401   |
| RT <sup>2</sup> SYBR Green ROX™ FAST Mastermix (2)* | For 2 x 96 assays in 96-well plates; suitable for use with the Rotor-Gene Q and other Rotor-Gene cyclers | 330620   |

\* Larger kit sizes available; please inquire.

RT<sup>2</sup> Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at [www.qiagen.com](http://www.qiagen.com) or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN<sup>®</sup>, Rotor-Gene<sup>®</sup>, Rotor-Disc™ (QIAGEN Group); ROX™ (Applied Biosystems or its subsidiaries); SYBR<sup>®</sup> (Molecular Probes, Inc.).

1067688 03/2011 © 2011 QIAGEN, all rights reserved.

[www.qiagen.com](http://www.qiagen.com)

Australia ■ 1-800-243-800

Austria ■ 0800/281010

Belgium ■ 0800-79612

Brazil ■ 0800-557779

Canada ■ 800-572-9613

China ■ 8621-3865-3865

Denmark ■ 80-885945

Finland ■ 0800-914416

France ■ 01-60-920-930

Germany ■ 02103-29-12000

Hong Kong ■ 800 933 965

Ireland ■ 1800 555 049

Italy ■ 800-787980

Japan ■ 03-6890-7300

Korea (South) ■ 080-000-7145

Luxembourg ■ 8002 2076

Mexico ■ 01-800-7742-436

The Netherlands ■ 0800 0229592

Norway ■ 800-18859

Singapore ■ 1800-742-4368

Spain ■ 91-630-7050

Sweden ■ 020-790282

Switzerland ■ 055-254-22-11

UK ■ 01293-422-911

USA ■ 800-426-8157



Sample & Assay Technologies